
# Geostationary Imaging FTS (GIFTS)

Breakthrough Technology
To Revolutionize the
Weather
Analysis/Forecast
Operation

W.L. Smith (NASA LaRC)

Weather Accident Prevention Annual Project Review

Hampton, Virginia, Radisson Hotel May 23-25, 2000






#### **Geostationary Imaging FTS (GIFTS)**



#### A Large Focal Plane Array Imaging FTS (Super-Sounder) for Geosynchronous Satellite Application



### Geostationary Imaging FTS (GIFTS)

#### Measurement Objectives:

## **Primary:** Greatly Improve Weather Forecasts By Measuring the Horizontal and Vertical Flux of Water Vapor

Observe Temperature, Water Vapor, Clouds, and Tracer Wind\* Profiles with HighVertical, Horizontal, and Temporal Resolution

## **Secondary:** Demonstrate Utility of Geostationary Satellites for Atmospheric Chemistry Studies


Observe Vertical Profiles and Transport of Radiatively Active Trace Gases including H<sub>2</sub>O, CO, O<sub>3</sub>, SO<sub>2</sub>

#### Technology Basis: Combine Measurement Technologies


Fourier Transform Spectrometer (FTS) with Large area format Focal Plane Array (LFPA) utilizing Active Cooling and On-board Digital Processing

\* Uses the Horizontal Displacement of Cloud and Retrieved Water Vapor Features


### GIFTS Sensor Module



## **GIFTS Spectral Coverage**




#### **GIFTS Sounding Accuracy**




#### **GIFTS Atmospheric Chemistry Measurements**


#### GIFTS averaging kernels for CO retrieval

#### GIFTS averaging kernels for ozone retrieval





#### Tropospheric Carbon Monoxide SRL-2 September 30 - October 11, 1994



## Geostationary Imaging FTS (GIFTS)

Measurement Capabilities: Imaging, Sounding, and Chemistry

Full Disk Imagery with 5 minute refresh rate, IR (4 Km) /VIS (1 Km)

Regional (6000 km x 6000 km) Imagery, Multi-spectral IR and Day/night Visible Imagery, with One (1) minute refresh rate

**Full Disk Sounding:** Hourly Temperature, Moisture, and Winds for Global Forecasting


Regional (6000 km x 6000 km) Sounding: Half Hourly, High Spatial Resolution Temperature, Moisture, and Wind Soundings for Regional Weather Forecasting

Mesoscale (3000 km x 3000 km) Sounding: Hourly, Ultra-high Vertical Resolution Soundings for Chemistry, Hazardous Weather Prediction Applications, and Self-Validation of Regional and Full Disk Products


## Airborne Demonstration GIFTS Measurement Capability




Wallops-98 (July 11, 1998) NAST-I Observations




#### NAST-I Demostrates GIFTS Water Vapor Profiling Capability (Sept. 14, 1998)





#### NAST-I Demonstration of GIFTS Wind Profiling Capability





## GIFTS Aviation Applications



- *Radiance Spectra* can be used to map cloud-top height, temperature, and ice/water phase (i.e., icing conditions), volcanic aerosol, fog, etc.
- *Soundings (T,q,V)* provide improved numerical weather forecasts and measure (i.e., nowcast) convective instability, wind shear, flight altitude winds, visibility, PBL height
- *Images* of derived products can be provided with a 5 30 minute frequency



## NASA and NAVY Partnership GIFTS/IOMI Enables Improved Global Weather Prediction









